

ON THE NATURE OF MATHEMATICAL TRUTH

C. G. HEMPEL, Queens College

1. The problem. It is a basic principle of scientific inquiry that no proposition and no theory is to be accepted without adequate grounds. In empirical science, which includes both the natural and the social sciences, the grounds for the acceptance of a theory consist in the agreement of predictions based on the theory with empirical evidence obtained either by experiment or by systematic observation. But what are the grounds which sanction the acceptance of mathematics? That is the question I propose to discuss in the present paper. For reasons which will become clear subsequently, I shall use the term "mathematics" here to refer to arithmetic, algebra, and analysis—to the exclusion, in particular, of geometry [1].

2. Are the propositions of mathematics self-evident truths? One of the several answers which have been given to our problem asserts that the truths of mathematics, in contradistinction to the hypotheses of empirical science, require neither factual evidence nor any other justification because they are "self-evident." This view, however, which ultimately relegates decisions as to mathematical truth to a feeling of self-evidence, encounters various difficulties. First of all, many mathematical theorems are so hard to establish that even to the specialist in the particular field they appear as anything but self-evident. Secondly, it is well known that some of the most interesting results of mathematics—especially in such fields as abstract set theory and topology—run counter to deeply ingrained intuitions and the customary kind of feeling of self-evidence. Thirdly, the existence of mathematical conjectures such as those of Goldbach and of Fermat, which are quite elementary in content and yet undecided up to this day, certainly shows that not all mathematical truths can be self-evident. And finally, even if self-evidence were attributed only to the basic postulates of mathematics, from which all other mathematical propositions can be deduced, it would be pertinent to remark that judgments as to what may be considered as self-evident are subjective; they may vary from person to person and certainly cannot constitute an adequate basis for decisions as to the objective validity of mathematical propositions.

3. Is mathematics the most general empirical science? According to another view, advocated especially by John Stuart Mill, mathematics is itself an empirical science which differs from the other branches such as astronomy, physics, chemistry, *etc.*, mainly in two respects: its subject matter is more general than that of any other field of scientific research, and its propositions have been tested and confirmed to a greater extent than those of even the most firmly established sections of astronomy or physics. Indeed, according to this view, the degree to which the laws of mathematics have been borne out by the past experiences of mankind is so overwhelming that—unjustifiably—we have come to think of mathematical theorems as qualitatively different from the well-confirmed hypotheses or theories of other branches of science: we consider them as

certain, while other theories are thought of as at best "very probable" or very highly confirmed.

But this view, too, is open to serious objections. From a hypothesis which is empirical in character—such as, for example, Newton's law of gravitation—it is possible to derive predictions to the effect that under certain specified conditions certain specified observable phenomena will occur. The actual occurrence of these phenomena constitutes confirming evidence, their non-occurrence disconfirming evidence for the hypothesis. It follows in particular that an empirical hypothesis is theoretically disconfirmable; *i.e.*, it is possible to indicate what kind of evidence, if actually encountered, would disconfirm the hypothesis. In the light of this remark, consider now a simple "hypothesis" from arithmetic: $3+2=5$. If this is actually an empirical generalization of past experiences, then it must be possible to state what kind of evidence would oblige us to concede the hypothesis was not generally true after all. If any disconfirming evidence for the given proposition can be thought of, the following illustration might well be typical of it: We place some microbes on a slide, putting down first three of them and then another two. Afterwards we count all the microbes to test whether in this instance 3 and 2 actually added up to 5. Suppose now that we counted 6 microbes altogether. Would we consider this as an empirical disconfirmation of the given proposition, or at least as a proof that it does not apply to microbes? Clearly not; rather, we would assume we had made a mistake in counting or that one of the microbes had split in two between the first and the second count. But under no circumstances could the phenomenon just described invalidate the arithmetical proposition in question; for the latter asserts nothing whatever about the behavior of microbes; it merely states that any set consisting of $3+2$ objects may also be said to consist of 5 objects. And this is so because the symbols " $3+2$ " and "5" denote the same number: they are synonymous by virtue of the fact that the symbols "2," "3," "5," and "+" are *defined* (or tacitly understood) in such a way that the above identity holds as a consequence of the meaning attached to the concepts involved in it.

4. The analytic character of mathematical propositions. The statement that $3+2=5$, then, is true for similar reasons as, say, the assertion that no sexagenarian is 45 years of age. Both are true simply by virtue of definitions or of similar stipulations which determine the meaning of the key terms involved. Statements of this kind share certain important characteristics: Their validation naturally requires no empirical evidence; they can be shown to be true by a mere analysis of the meaning attached to the terms which occur in them. In the language of logic, sentences of this kind are called analytic or true a priori, which is to indicate that their truth is logically independent of, or logically prior to, any experiential evidence [2]. And while the statements of empirical science, which are synthetic and can be validated only a posteriori, are constantly subject to revision in the light of new evidence, the truth of an analytic statement can be established definitely, once and for all. However, this characteristic "theoretical

certainty" of analytic propositions has to be paid for at a high price: An analytic statement conveys no factual information. Our statement about sexagenarians, for example, asserts nothing that could possibly conflict with any factual evidence: it has no factual implications, no empirical content; and it is precisely for this reason that the statement can be validated without recourse to empirical evidence.

Let us illustrate this view of the nature of mathematical propositions by reference to another, frequently cited, example of a mathematical—or rather logical—truth, namely the proposition that whenever $a = b$ and $b = c$ then $a = c$. On what grounds can this so-called "transitivity of identity" be asserted? Is it of an empirical nature and hence at least theoretically disconfirmable by empirical evidence? Suppose, for example, that a, b, c , are certain shades of green, and that as far as we can see, $a = b$ and $b = c$, but clearly $a \neq c$. This phenomenon actually occurs under certain conditions; do we consider it as disconfirming evidence for the proposition under consideration? Undoubtedly not; we would argue that if $a \neq c$, it is impossible that $a = b$ and also $b = c$; between the terms of at least one of these latter pairs, there must obtain a difference, though perhaps only a subliminal one. And we would dismiss the possibility of empirical disconfirmation, and indeed the idea that an empirical test should be relevant here, on the grounds that identity is a transitive relation by virtue of its definition or by virtue of the basic postulates governing it [3]. Hence, the principle in question is true a priori.

5. Mathematics as an axiomatized deductive system. I have argued so far that the validity of mathematics rests neither on its alleged self-evidential character nor on any empirical basis, but derives from the stipulations which determine the meaning of the mathematical concepts, and that the propositions of mathematics are therefore essentially "true by definition." This latter statement, however, is obviously oversimplified and needs restatement and a more careful justification.

For the rigorous development of a mathematical theory proceeds not simply from a set of definitions but rather from a set of non-definitional propositions which are not proved within the theory; these are the postulates or axioms of the theory [4]. They are formulated in terms of certain basic or primitive concepts for which no definitions are provided within the theory. It is sometimes asserted that the postulates themselves represent "implicit definitions" of the primitive terms. Such a characterization of the postulates, however, is misleading. For while the postulates do limit, in a specific sense, the meanings that can possibly be ascribed to the primitives, any self-consistent postulate system admits, nevertheless, many different interpretations of the primitive terms (this will soon be illustrated), whereas a set of definitions in the strict sense of the word determines the meanings of the definienda in a unique fashion.

Once the primitive terms and the postulates have been laid down, the entire theory is completely determined; it is derivable from its postulational basis in

the following sense: Every term of the theory is definable in terms of the primitives, and every proposition of the theory is logically deducible from the postulates. To be entirely precise, it is necessary also to specify the principles of logic which are to be used in the proof of the propositions, *i.e.* in their deduction from the postulates. These principles can be stated quite explicitly. They fall into two groups: Primitive sentences, or postulates, of logic (such as: If p and q is the case, then p is the case), and rules of deduction or inference (including, for example, the familiar modus ponens rule and the rules of substitution which make it possible to infer, from a general proposition, any one of its substitution instances). A more detailed discussion of the structure and content of logic would, however, lead too far afield in the context of this article.

6. Peano's axiom system as a basis for mathematics. Let us now consider a postulate system from which the entire arithmetic of the natural numbers can be derived. This system was devised by the Italian mathematician and logician G. Peano (1858–1932). The primitives of this system are the terms “0,” “number,” and “successor.” While, of course, no definition of these terms is given within the theory, the symbol “0” is intended to designate the number 0 in its usual meaning, while the term “number” is meant to refer to the natural numbers 0, 1, 2, 3 . . . exclusively. By the successor of a natural number n , which will sometimes briefly be called n' , is meant the natural number immediately following n in the natural order. Peano's system contains the following 5 postulates:

- P1. 0 is a number
- P2. The successor of any number is a number
- P3. No two numbers have the same successor
- P4. 0 is not the successor of any number
- P5. If P is a property such that (a) 0 has the property P , and (b) whenever a number n has the property P , then the successor of n also has the property P , then every number has the property P .

The last postulate embodies the principle of mathematical induction and illustrates in a very obvious manner the enforcement of a mathematical “truth” by stipulation. The construction of elementary arithmetic on this basis begins with the definition of the various natural numbers. 1 is defined as the successor of 0, or briefly as $0'$; 2 as $1'$, 3 as $2'$, and so on. By virtue of P2, this process can be continued indefinitely; because of P3 (in combination with P5), it never leads back to one of the numbers previously defined, and in view of P4, it does not lead back to 0 either.

As the next step, we can set up a definition of addition which expresses in a precise form the idea that the addition of any natural number to some given number may be considered as a repeated addition of 1; the latter operation is readily expressible by means of the successor relation. This definition of addition runs as follows:

D1. (a) $n+0=n$; (b) $n+k'=(n+k)'$.

The two stipulations of this recursive definition completely determine the sum of any two integers. Consider, for example, the sum $3+2$. According to the definitions of the numbers 2 and 1, we have $3+2=3+1'=3+(0)'$; by D1 (b), $3+(0)'=(3+0)'=((3+0)')'$; but by D1 (a), and by the definitions of the numbers 4 and 5, $((3+0)')'=(3')'=4'=5$. This proof also renders more explicit and precise the comments made earlier in this paper on the truth of the proposition that $3+2=5$: Within the Peano system of arithmetic, its truth flows not merely from the definition of the concepts involved, but also from the postulates that govern these various concepts. (In our specific example, the postulates P1 and P2 are presupposed to guarantee that 1, 2, 3, 4, 5 are numbers in Peano's system; the general proof that D1 determines the sum of any two numbers also makes use of P5.) If we call the postulates and definitions of an axiomatized theory the "stipulations" concerning the concepts of that theory, then we may say now that the propositions of the arithmetic of the natural numbers are true by virtue of the stipulations which have been laid down initially for the arithmetical concepts. (Note, incidentally, that our proof of the formula " $3+2=5$ " repeatedly made use of the transitivity of identity; the latter is accepted here as one of the rules of logic which may be used in the proof of any arithmetical theorem; it is, therefore, included among Peano's postulates no more than any other principle of logic.)

Now, the multiplication of natural numbers may be defined by means of the following recursive definition, which expresses in a rigorous form the idea that a product nk of two integers may be considered as the sum of k terms each of which equals n .

D2. (a) $n \cdot 0 = 0$; (b) $n \cdot k' = n \cdot k + n$.

It now is possible to prove the familiar general laws governing addition and multiplication, such as the commutative, associative, and distributive laws ($n+k=k+n$, $n \cdot k=k \cdot n$; $n+(k+l)=(n+k)+l$, $n \cdot (k \cdot l)=(n \cdot k) \cdot l$; $n \cdot (k+l)=(n \cdot k)+(n \cdot l)$).—In terms of addition and multiplication, the inverse operations of subtraction and division can then be defined. But it turns out that these "cannot always be performed"; *i.e.*, in contradistinction to the sum and the product, the difference and the quotient are not defined for every couple of numbers; for example, $7-10$ and $7 \div 10$ are undefined. This situation suggests an enlargement of the number system by the introduction of negative and of rational numbers.

It is sometimes held that in order to effect this enlargement, we have to "assume" or else to "postulate" the existence of the desired additional kinds of numbers with properties that make them fit to fill the gaps of subtraction and division. This method of simply postulating what we want has its advantages; but, as Bertrand Russell [5] puts it, they are the same as the advantages of theft over honest toil; and it is a remarkable fact that the negative as well as the ra-

tional numbers can be obtained from Peano's primitives by the honest toil of constructing explicit definitions for them, without the introduction of any new postulates or assumptions whatsoever. Every positive and negative integer—in contradistinction to a natural number which has no sign—is definable as a certain set of ordered couples of natural numbers; thus, the integer $+2$ is definable as the set of all ordered couples (m, n) of natural numbers where $m = n + 2$; the integer -2 is the set of all ordered couples (m, n) of natural numbers with $n = m + 2$.—Similarly, rational numbers are defined as classes of ordered couples of integers.—The various arithmetical operations can then be defined with reference to these new types of numbers, and the validity of all the arithmetical laws governing these operations can be proved by virtue of nothing more than Peano's postulates and the definitions of the various arithmetical concepts involved.

The much broader system thus obtained is still incomplete in the sense that not every number in it has a square root, and more generally, not every algebraic equation whose coefficients are all numbers of the system has a solution in the system. This suggests further expansions of the number system by the introduction of real and finally of complex numbers. Again, this enormous extension can be effected by mere definition, without the introduction of a single new postulate [6]. On the basis thus obtained, the various arithmetical and algebraic operations can be defined for the numbers of the new system, the concepts of function, of limit, of derivative and integral can be introduced, and the familiar theorems pertaining to these concepts can be proved, so that finally the huge system of mathematics as here delimited rests on the narrow basis of Peano's system: Every concept of mathematics can be defined by means of Peano's three primitives, and every proposition of mathematics can be deduced from the five postulates enriched by the definitions of the non-primitive terms [6a]. These deductions can be carried out, in most cases, by means of nothing more than the principles of formal logic; the proof of some theorems concerning real numbers, however, requires one assumption which is not usually included among the latter. This is the so-called axiom of choice. It asserts that given a class of mutually exclusive classes, none of which is empty, there exists at least one class which has exactly one element in common with each of the given classes. By virtue of this principle and the rules of formal logic, the content of all of mathematics can thus be derived from Peano's modest system—a remarkable achievement in systematizing the content of mathematics and clarifying the foundations of its validity.

7. Interpretations of Peano's primitives. As a consequence of this result, the whole system of mathematics might be said to be true by virtue of mere definitions (namely, of the non-primitive mathematical terms) provided that the five Peano postulates are true. However, strictly speaking, we cannot, at this juncture, refer to the Peano postulates as propositions which are either true or false, for they contain three primitive terms which have not been assigned any specific meaning. All we can assert so far is that any specific interpretation of the primi-

tives which satisfies the five postulates—*i.e.*, turns them into true statements—will also satisfy all the theorems deduced from them. But for Peano's system, there are several—indeed, infinitely many—interpretations which will do this. For example, let us understand by 0 the origin of a half-line, by the successor of a point on that half-line the point 1 cm. behind it, counting from the origin, and by a number any point which is either the origin or can be reached from it by a finite succession of steps each of which leads from one point to its successor. It can then readily be seen that all the Peano postulates as well as the ensuing theorems turn into true propositions, although the interpretation given to the primitives is certainly not the customary one, which was mentioned earlier. More generally, it can be shown that every progression of elements of any kind provides a true interpretation, or a "model," of the Peano system. This example illustrates our earlier observation that a postulate system cannot be regarded as a set of "implicit definitions" for the primitive terms: The Peano system permits of many different interpretations, whereas in everyday as well as in scientific language, we attach one specific meaning to the concepts of arithmetic. Thus, *e.g.*, in scientific and in everyday discourse, the concept 2 is understood in such a way that from the statement "Mr. Brown as well as Mr. Cope, but no one else is in the office, and Mr. Brown is not the same person as Mr. Cope," the conclusion "Exactly two persons are in the office" may be validly inferred. But the stipulations laid down in Peano's system for the natural numbers, and for the number 2 in particular, do not enable us to draw this conclusion; they do not "implicitly determine" the customary meaning of the concept 2 or of the other arithmetical concepts. And the mathematician cannot acquiesce at this deficiency by arguing that he is not concerned with the customary meaning of the mathematical concepts; for in proving, say, that every positive real number has exactly two real square roots, he is himself using the concept 2 in its customary meaning, and his very theorem cannot be proved unless we presuppose more about the number 2 than is stipulated in the Peano system.

If therefore mathematics is to be a correct theory of the mathematical concepts in their intended meaning, it is not sufficient for its validation to have shown that the entire system is derivable from the Peano postulates plus suitable definitions; rather, we have to inquire further whether the Peano postulates are actually true when the primitives are understood in their customary meaning. This question, of course, can be answered only after the customary meaning of the terms "0," "natural number," and "successor" has been clearly defined. To this task we now turn.

8. Definition of the customary meaning of the concepts of arithmetic in purely logical terms. At first blush, it might seem a hopeless undertaking to try to define these basic arithmetical concepts without presupposing other terms of arithmetic, which would involve us in a circular procedure. However, quite rigorous definitions of the desired kind can indeed be formulated, and it can be shown that for the concepts so defined, all Peano postulates turn into true state-

ments. This important result is due to the research of the German logician G. Frege (1848–1925) and to the subsequent systematic and detailed work of the contemporary English logicians and philosophers B. Russell and A. N. Whitehead. Let us consider briefly the basic ideas underlying these definitions [7].

A natural number—or, in Peano's term, a number—in its customary meaning can be considered as a characteristic of certain *classes* of objects. Thus, *e.g.*, the class of the apostles has the number 12, the class of the Dionne quintuplets the number 5, any couple the number 2, and so on. Let us now express precisely the meaning of the assertion that a certain class C has the number 2, or briefly, that $n(C) = 2$. Brief reflection will show that the following definiens is adequate in the sense of the customary meaning of the concept 2: There is some object x and some object y such that (1) $x \in C$ (*i.e.*, x is an element of C) and $y \in C$, (2) $x \neq y$, and (3) if z is any object such that $z \in C$, then either $z = x$ or $z = y$. (Note that on the basis of this definition it becomes indeed possible to infer the statement "The number of persons in the office is 2" from "Mr. Brown as well as Mr. Cope, but no one else is in the office, and Mr. Brown is not identical with Mr. Cope"; C is here the class of persons in the office.) Analogously, the meaning of the statement that $n(C) = 1$ can be defined thus: There is some x such that $x \in C$, and any object y such that $y \in C$, is identical with x . Similarly, the customary meaning of the statement that $n(C) = 0$ is this: There is no object such that $x \in C$.

The general pattern of these definitions clearly lends itself to the definition of any natural number. Let us note especially that in the definitions thus obtained, the definiens never contains any arithmetical term, but merely expressions taken from the field of formal logic, including the signs of identity and difference. So far, we have defined only the meaning of such phrases as " $n(C) = 2$," but we have given no definition for the numbers 0, 1, 2, . . . apart from this context. This desideratum can be met on the basis of the consideration that 2 is that property which is common to all couples, *i.e.*, to all classes C such that $n(C) = 2$. This common property may be conceptually represented by the class of all those classes which share this property. Thus we arrive at the definition: 2 is the class of all couples, *i.e.*, the class of all classes C for which $n(C) = 2$.—This definition is by no means circular because the concept of couple—in other words, the meaning of " $n(C) = 2$ "—has been previously defined without any reference to the number 2. Analogously, 1 is the class of all unit classes, *i.e.*, the class of all classes C for which $n(C) = 1$. Finally, 0 is the class of all null classes, *i.e.*, the class of all classes without elements. And as there is only one such class, 0 is simply the class whose only element is the null class. Clearly, the customary meaning of any given natural number can be defined in this fashion [8]. In order to characterize the intended interpretation of Peano's primitives, we actually need, of all the definitions here referred to, only that of the number 0. It remains to define the terms "successor" and "integer."

The definition of "successor," whose precise formulation involves too many

niceties to be stated here, is a careful expression of a simple idea which is illustrated by the following example: Consider the number 5, *i.e.*, the class of all quintuplets. Let us select an arbitrary one of these quintuplets and add to it an object which is not yet one of its members. $5'$, the successor of 5, may then be defined as the number applying to the set thus obtained (which, of course, is a sextuplet). Finally, it is possible to formulate a definition of the customary meaning of the concept of natural number; this definition, which again cannot be given here, expresses, in a rigorous form, the idea that the class of the natural numbers consists of the number 0, its successor, the successor of that successor, and so on.

If the definitions here characterized are carefully written out—this is one of the cases where the techniques of symbolic, or mathematical, logic prove indispensable—it is seen that the definiens of every one of them contains exclusively terms from the field of pure logic. In fact, it is possible to state the customary interpretation of Peano's primitives, and thus also the meaning of every concept definable by means of them—and that includes every concept of mathematics—in terms of the following 7 expressions, in addition to variables such as " x " and " C ": *not*, *and*, *if—then*; *for every object x it is the case that . . .*; *there is some object x such that . . .*; *x is an element of class C* ; *the class of all things x such that . . .*. And it is even possible to reduce the number of logical concepts needed to a mere four: The first three of the concepts just mentioned are all definable in terms of "*neither—nor*," and the fifth is definable by means of the fourth and "*neither—nor*." Thus, all the concepts of mathematics prove definable in terms of four concepts of pure logic. (The definition of one of the more complex concepts of mathematics in terms of the four primitives just mentioned may well fill hundreds or even thousands of pages; but clearly this affects in no way the theoretical importance of the result just obtained; it does, however, show the great convenience and indeed practical indispensability for mathematics of having a large system of highly complex defined concepts available.)

9. The truth of Peano's postulates in their customary interpretation. The definitions characterized in the preceding section may be said to render precise and explicit the customary meaning of the concepts of arithmetic. Moreover—and this is crucial for the question of the validity of mathematics—it can be shown that the Peano postulates all turn into true propositions if the primitives are construed in accordance with the definitions just considered.

Thus, P1 (0 is a number) is true because the class of all numbers—*i.e.*, natural numbers—was defined as consisting of 0 and all its successors. The truth of P2 (The successor of any number is a number) follows from the same definition. This is true also of P5, the principle of mathematical induction. To prove this, however, we would have to resort to the precise definition of "integer" rather than the loose description given of that definition above. P4 (0 is not the successor of any number) is seen to be true as follows: By virtue of the definition of "successor," a number which is a successor of some number can apply only to

classes which contain at least one element; but the number 0, by definition, applies to a class if and only if that class is empty.—While the truth of P1, P2, P4, P5 can be inferred from the above definitions simply by means of the principles of logic, the proof of P3 (No two numbers have the same successor) presents a certain difficulty. As was mentioned in the preceding section, the definition of the successor of a number n is based on the process of adding, to a class of n elements, one element not yet contained in that class. Now if there should exist only a finite number of things altogether then this process could not be continued indefinitely, and P3, which (in conjunction with P1 and P2) implies that the integers form an infinite set, would be false. Russell's way of meeting this difficulty [9] was to introduce a special "axiom of infinity," which stipulates, in effect, the existence of infinitely many objects and thus makes P3 demonstrable. The axiom of infinity can be formulated in purely logical terms and may therefore be considered as a postulate of logic; however, it certainly does not belong to the generally recognized principles of logic; and it thus introduces a foreign element into the otherwise unexceptionable derivation of the Peano postulates from pure logic. Recently, however, it has been shown [10] that a suitable system of logical principles can be set up which is even less comprehensive than the rules of logic which are commonly used [11], and in which the existence of infinitely many objects can be proved without the need for a special axiom.

10. Mathematics as a branch of logic. As was pointed out earlier, all the theorems of arithmetic, algebra, and analysis can be deduced from the Peano postulates and the definitions of those mathematical terms which are not primitives in Peano's system. This deduction requires only the principles of logic plus, in certain cases, the axiom of choice. By combining this result with what has just been said about the Peano system, the following conclusion is obtained, which is also known as *the thesis of logicism concerning the nature of mathematics*:

Mathematics is a branch of logic. It can be derived from logic in the following sense:

- a. All the concepts of mathematics, *i.e.* of arithmetic, algebra, and analysis, can be defined in terms of four concepts of pure logic.
- b. All the theorems of mathematics can be deduced from those definitions by means of the principles of logic (including the axiom of choice).

In this sense it can be said that the propositions of the system of mathematics as here delimited are true by virtue of the definitions of the mathematical concepts involved, or that they make explicit certain characteristics with which we have endowed our mathematical concepts by definition. The propositions of mathematics have, therefore, the same unquestionable certainty which is typical of such propositions as "All bachelors are unmarried," but they also share the complete lack of empirical content which is associated with that certainty: The propositions of mathematics are devoid of all factual content; they convey no information whatever on any empirical subject matter.

11. On the applicability of mathematics to empirical subject matter. This result seems to be irreconcilable with the fact that after all mathematics has proved to be eminently applicable to empirical subject matter, and that indeed the greater part of present-day scientific knowledge has been reached only through continual reliance on and application of the propositions of mathematics.—Let us try to clarify this apparent paradox by reference to some examples.

Suppose that we are examining a certain amount of some gas, whose volume v , at a certain fixed temperature, is found to be 9 cubic feet when the pressure p is 4 atmospheres. And let us assume further that the volume of the gas for the same temperature and $p=6$ *at.*, is predicted by means of Boyle's law. Using elementary arithmetic we reason thus: For corresponding values of v and p , $vp=c$, and $v=9$ when $p=4$; hence $c=36$: Therefore, when $p=6$, then $v=6$. Suppose that this prediction is borne out by subsequent test. Does that show that the arithmetic used has a predictive power of its own, that its propositions have factual implications? Certainly not. All the predictive power here deployed, all the empirical content exhibited stems from the initial data and from Boyle's law, which asserts that $vp=c$ for *any* two corresponding values of v and p , hence also for $v=9$, $p=4$, and for $p=6$ and the corresponding value of v [12]. The function of the mathematics here applied is not predictive at all; rather, it is analytic or explicative: it renders explicit certain assumptions or assertions which are included in the content of the premises of the argument (in our case, these consist of Boyle's law plus the additional data); mathematical reasoning reveals that those premises contain—hidden in them,—as it were,—an assertion about the case as yet unobserved. In accepting our premises—so arithmetic reveals—we have—knowingly or unknowingly—already accepted the implication that the p -value in question is 6. Mathematical as well as logical reasoning is a conceptual technique of making explicit what is implicitly contained in a set of premises. The conclusions to which this technique leads assert nothing that is *theoretically new* in the sense of not being contained in the content of the premises. But the results obtained may well be *psychologically new*: we may not have been aware, before using the techniques of logic and mathematics, what we committed ourselves to in accepting a certain set of assumptions or assertions.

A similar analysis is possible in all other cases of applied mathematics, including those involving, say, the calculus. Consider, for example, the hypothesis that a certain object, moving in a specified electric field, will undergo a constant acceleration of 5 feet/sec². For the purpose of testing this hypothesis, we might derive from it, by means of two successive integrations, the prediction that if the object is at rest at the beginning of the motion, then the distance covered by it at any time t is $\frac{5}{2}t^2$ feet. This conclusion may clearly be psychologically new to a person not acquainted with the subject, but it is not theoretically new; the content of the conclusion is already contained in that of the hypothesis about the constant acceleration. And indeed, here as well as in the case of the compression of a gas, a failure of the prediction to come true would be considered as indica-

tive of the factual incorrectness of at least one of the premises involved (*f.ex.*, of Boyle's law in its application to the particular gas), but never as a sign that the logical and mathematical principles involved might be unsound.

Thus, in the establishment of empirical knowledge, mathematics (as well as logic) has, so to speak, the function of a theoretical juice extractor: the techniques of mathematical and logical theory can produce no more juice of factual information than is contained in the assumptions to which they are applied; but they may produce a great deal more juice of this kind than might have been anticipated upon a first intuitive inspection of those assumptions which form the raw material for the extractor.

At this point, it may be well to consider briefly the status of those mathematical disciplines which are not outgrowths of arithmetic and thus of logic; these include in particular topology, geometry, and the various branches of abstract algebra, such as the theory of groups, lattices, fields, *etc.* Each of these disciplines can be developed as a purely deductive system on the basis of a suitable set of postulates. If P be the conjunction of the postulates for a given theory, then the proof of a proposition T of that theory consists in deducing T from P by means of the principles of formal logic. What is established by the proof is therefore not the truth of T , but rather the fact that T is true provided that the postulates are. But since both P and T contain certain primitive terms of the theory, to which no specific meaning is assigned, it is not strictly possible to speak of the truth of either P or T ; it is therefore more adequate to state the point as follows: If a proposition T is logically deduced from P , then every specific interpretation of the primitives which turns all the postulates of P into true statements, will also render T a true statement.—Up to this point, the analysis is exactly analogous to that of arithmetic as based on Peano's set of postulates. In the case of arithmetic, however, it proved possible to go a step further, namely to define the customary meanings of the primitives in terms of purely logical concepts and to show that the postulates—and therefore also the theorems—of arithmetic are unconditionally true by virtue of these definitions. An analogous procedure is not applicable to those disciplines which are not outgrowths of arithmetic: The primitives of the various branches of abstract algebra have no specific "customary meaning"; and if geometry in its customary interpretation is thought of as a theory of the structure of physical space, then its primitives have to be construed as referring to certain types of physical entities, and the question of the truth of a geometrical theory in this interpretation turns into an *empirical* problem [13]. For the purpose of applying any one of these non-arithmetical disciplines to some specific field of mathematics or empirical science, it is therefore necessary first to assign to the primitives some specific meaning and then to ascertain whether in this interpretation the postulates turn into true statements. If this is the case, then we can be sure that all the theorems are true statements too, because they are logically derived from the postulates and thus simply explicate the content of the latter in the given interpretation.—In their application to empirical subject matter, therefore, these mathematical theories no less than

those which grow out of arithmetic and ultimately out of pure logic, have the function of an analytic tool, which brings to light the implications of a given set of assumptions but adds nothing to their content.

But while mathematics in no case contributes anything to the content of our knowledge of empirical matters, it is entirely indispensable as an instrument for the validation and even for the linguistic expression of such knowledge: The majority of the more far-reaching theories in empirical science—including those which lend themselves most eminently to prediction or to practical application—are stated with the help of mathematical concepts; the formulation of these theories makes use, in particular, of the number system, and of functional relationships among different metrical variables. Furthermore, the scientific test of these theories, the establishment of predictions by means of them, and finally their practical application, all require the deduction, from the general theory, of certain specific consequences; and such deduction would be entirely impossible without the techniques of mathematics which reveal what the given general theory implicitly asserts about a certain special case.

Thus, the analysis outlined on these pages exhibits the system of mathematics as a vast and ingenious conceptual structure without empirical content and yet an indispensable and powerful theoretical instrument for the scientific understanding and mastery of the world of our experience.

References

1. A discussion of the status of geometry is given in my article, *Geometry and Empirical Science*, *American Mathematical Monthly*, vol. 52, pp. 7-17; 1945.
2. The objection is sometimes raised that without certain types of experience, such as encountering several objects of the same kind, the integers and the arithmetical operations with them would never have been invented, and that therefore the propositions of arithmetic do have an empirical basis. This type of argument, however, involves a confusion of the logical and the psychological meaning of the term "basis." It may very well be the case that certain experiences occasion psychologically the formation of arithmetical ideas and in this sense form an empirical "basis" for them; but this point is entirely irrelevant for the logical questions as to the *grounds* on which the propositions of arithmetic may be accepted as true. The point made above is that no empirical "basis" or evidence whatever is needed to establish the truth of the propositions of arithmetic.
3. A precise account of the definition and the essential characteristics of the identity relation may be found in A. Tarski, *Introduction to Logic*, New York, 1941, Ch. III.
4. For a lucid and concise account of the axiomatic method, see A. Tarski, *l.c.*, Ch. VI.
5. Bertrand Russell, *Introduction to Mathematical Philosophy*, New York and London, 1919, p. 71.
6. For a more detailed account of the construction of the number system on Peano's basis, cf. Bertrand Russell, *l.c.*, esp. Chs. I and VII.—A rigorous and concise presentation of that construction, beginning, however, with the set of all integers rather than that of the natural numbers, may be found in G. Birkhoff and S. MacLane, *A Survey of Modern Algebra*, New York 1941, Chs. I, II, III, V.—For a general survey of the construction of the number system, cf. also J. W. Young, *Lectures on the Fundamental Concepts of Algebra and Geometry*, New York, 1911, esp. lectures X, XI, XII.
- 6a. As a result of very deep-reaching investigations carried out by K. Gödel it is known that arithmetic, and *a fortiori* mathematics, is an incomplete theory in the following sense: While all those propositions which belong to the classical systems of arithmetic, algebra, and analysis can

indeed be derived, in the sense characterized above, from the Peano postulates, there exist nevertheless other propositions which can be expressed in purely arithmetical terms, and which are true, but which cannot be derived from the Peano system. And more generally: For any postulate system of arithmetic (or of mathematics for that matter) which is not self-contradictory, there exist propositions which are true, and which can be stated in purely arithmetical terms, but which cannot be derived from that postulate system. In other words, it is impossible to construct a postulate system which is not self-contradictory, and which contains among its consequences all true propositions which can be formulated within the language of arithmetic.

This fact does not, however, affect the result outlined above, namely, that it is possible to deduce, from the Peano postulates and the additional definitions of non-primitive terms, all those propositions which constitute the classical theory of arithmetic, algebra, and analysis; and it is to these propositions that I refer above and subsequently as the propositions of mathematics.

7. For a more detailed discussion, *cf.* Russell, *l.c.*, Chs. II, III, IV. A complete technical development of the idea can be found in the great standard work in mathematical logic, A. N. Whitehead and B. Russell, *Principia Mathematica*, Cambridge, England, 1910–1913.—For a very precise recent development of the theory, see W. V. O. Quine, *Mathematical Logic*, New York 1940.—A specific discussion of the Peano system and its interpretations from the viewpoint of semantics is included in R. Carnap, *Foundations of Logic and Mathematics*, *International Encyclopedia of Unified Science*, vol. I, no. 3, Chicago, 1939; especially sections 14, 17, 18.

8. The assertion that the definitions given above state the “customary” meaning of the arithmetical terms involved is to be understood in the logical, not the psychological sense of the term “meaning.” It would obviously be absurd to claim that the above definitions express “what everybody has in mind” when talking about numbers and the various operations that can be performed with them. What is achieved by those definitions is rather a “logical reconstruction” of the concepts of arithmetic in the sense that if the definitions are accepted, then those statements in science and everyday discourse which involve arithmetical terms can be interpreted coherently and systematically in such a manner that they are capable of objective validation. The statement about the two persons in the office provides a very elementary illustration of what is meant here.

9. *Cf.* Bertrand Russell, *l.c.*, p. 24 and Ch. XIII.

10. This result has been obtained by W. V. O. Quine; *cf.* his *Mathematical Logic*, New York, 1940.

11. The principles of logic developed in Quine's work and in similar modern systems of formal logic embody certain restrictions as compared with those logical rules which had been rather generally accepted as sound until about the turn of the 20th century. At that time, the discovery of the famous paradoxes of logic, especially of Russell's paradox (*cf.* Russell, *l.c.*, Ch. XIII) revealed the fact that the logical principles implicit in customary mathematical reasoning involved contradictions and therefore had to be curtailed in one manner or another.

12. Note that we may say “hence” by virtue of the rule of substitution, which is one of the rules of logical inference.

13. For a more detailed discussion of this point, *cf.* the article mentioned in reference 1.